

GALACTIC WINDS IN IRREGULAR STARBURST GALAXIES

F. MATTEUCCI

*Department of Astronomy, Trieste University, Via G.B. Tiepolo 11,
34100 Trieste, Italy
E-mail: matteucci@ts.astro.it*

S. RECCHI

*Department of Astronomy, Trieste University, Via G.B. Tiepolo 11,
34100 Trieste, Italy
E-mail: recchi@ts.astro.it*

In this paper we present some results concerning the study of the development of galactic winds in blue compact galaxies. In particular, we model a situation very similar to that of the galaxy IZw18, the most metal poor and unevolved galaxy known locally. To do that we compute the chemo-dynamical evolution of a galaxy in the case of one instantaneous isolated starburst as well as in the case of two successive instantaneous starbursts. We show that in both cases a metal enriched wind develops and that the metals produced by the type Ia SNe are lost more efficiently than those produced by type II SNe. We also find that one single burst is able to enrich chemically the surrounding region in few Myr. Both these results are the effect of the assumed efficiency of energy transfer from SNe to ISM and to the consideration of type Ia SNe in this kind of problem. The comparison with observed abundances of IZw18 suggests that this galaxy is likely to have suffered two bursts in its life, with the previous being less intense than the last one.

1 One and two bursts of star formation

We assume a single instantaneous burst occurring at the center of a gas-rich dwarf surrounded by a dark matter halo. The mass of stars formed is $M_* = 6 \cdot 10^5 M_\odot$, the gas mass is $M_{gas} = 1.7 \cdot 10^7 M_\odot$ and the mass of the dark halo is $M_{dark} = 6.5 \cdot 10^8 M_\odot$. These parameters are chosen to reproduce, for what possible, the characteristics of IZw18 (unevolved local galaxy). The galactic region extends for 700 pc in the vertical (z) direction and for 1kpc in the radial direction (R). In order to study the chemo-dynamical evolution consequent to the burst, we adopt a 2-D hydrocode coupled with detailed chemical yields from: type II SNe, type Ia SNe and low and intermediate mass stars (see Recchi et al. 2001 for more details about this model). We follow the evolution of the abundances of H, He, C, N, O, Mg, Si, Fe in the gas. The evolution is followed for 375 Myr since the burst. We also consider a case with a second burst occurring after 300 Myr from the first one. The mass in stars formed during the first burst is $M_* = 10^5 M_\odot$, whereas the mass in stars formed in the second burst is $M_* = 5.8 \cdot 10^5 M_\odot$. The initial gas mass and dark matter

halo are the same as in the one-burst case. The simulation lasts 450 Myr in total since the first burst. The SN efficiencies of energy transfer into the ISM from SNe is the same in both cases, in particular: for type II SNe we assume that the efficiency is $\eta = 0.03$ and for type Ia SNe $\eta = 1$. This choice is due to results from Bradamante et al. (1998) for type II SNe indicating that the first SNe to explode lose a large fraction of their initial energy by radiation due to the cold and dense ISM in their surroundings, whereas for type Ia SNe the efficiency is maximum due to the fact that they explode into an ISM already hot and rarefied (see Recchi et al. 2001, for details).

2 Dynamical Results

We find that the starburst triggers indeed a galactic wind and the metals leave the galaxy more easily than the unprocessed gas confirming previous results (e.g. McLow and Ferrara, 1999). We find that SNe Ia eject their metals more efficiently than SNe II since they inject all of their initial energy into the ISM. This is a new result relative to previous studies since it is the first time that type Ia SNe are taken into account. At variance with previous studies (see e.g. Tenorio-Tagle 1996) we find that most of the metals are already in the cold gas phase after 8-10 Myr from the beginning of the burst, due to the fact that the superbubble created by the SNe does not break immediately and thermal conduction can act efficiently.

3 Chemical results

We find that one single instantaneous burst, occurring in a primordial gas (no metallicity), at an age of ~ 31 Myr can reasonably reproduce the abundances measured in IZw18 (see figure 1). From this one would conclude that perhaps this galaxy is experiencing its first burst of SF, although one cannot exclude a previous burst which enriched the gas no more than $1/50 Z_{\odot}$. However, as evident in figure 1, the correct N/O ratio would last only for a very short time since for $t > 31$ Myr the N/O ratio will start to increase outside the permitted observational range. In addition, Color Magnitude Diagram studies (e.g. Aloisi et al. 1999) indicate the presence of an old underlying stellar population in IZw18, thus suggesting that the two-burst case is more realistic. An interesting result is that the $[\alpha/\text{Fe}]$ ratios in the gas outside the galaxy (i.e. in the galactic wind) are lower than inside, due to the larger ejection efficiency by type Ia SNe (more iron is lost than α -elements). This creates an interesting dichotomy which can have important consequences on the evolution of the ICM, especially if this effect will be found also in elliptical galaxies.




Figure 1: Evolution of O, C, N for the single-burst model (left panels) and the double-burst model (right panels). The superimposed grids represent the observative values found in literature for IZw18

In the two-burst case, at the onset of the second burst the metallicity in the burst region is $\sim Z_{\odot}/50$, and a very good agreement with the measured abundances in IZw18 is obtained in this case, especially for the C/O and N/O ratios and for a larger range of burst ages (see figure 1). Our results suggest that a first weak burst occurred more than 300 Myr ago and followed by a more intense one with an age between 20 and 70 Myr can reproduce most of the properties of this galaxy. The $[\alpha/\text{Fe}]$ ratios in the two burst case are always lower outside than inside the galaxy, but the effect is less evident than in the one-burst case.

Acknowledgments

We are grateful to Annibale D’Ercole and Monica Tosi for their help in this work.

References

1. Aloisi A., Tosi M. and Greggio L., *AJ* **118**, 302 (1999).
2. Bradamante F., Matteucci F. and D’Ercole A., *A&A* **337**, 338 (1998).
3. Mac Low M.-M. and Ferrara A., *ApJ* **513**, 142 (1999).
4. Recchi S., Matteucci F. and D’Ercole A., *MNRAS* in press (2001).
5. Tenorio-Tagle G., *AJ* **111**, 1641 (1996)